



Improving Patient-Provider Communication

A Call to Action

Lance Patak, MD, MBA, RN
Amy Wilson-Stronks, MPP, CPHQ
John Costello, MA, CCC-SLP
Ruth M. Kleinpell, PhD, RN, FAAN
Elizabeth A. Henneman, PhD, RN
Colleen Person, MMA, BSN, RN
Mary Beth Happ, PhD, RN

Patients who are communication impaired are at greater risk of medical error and poorer outcomes. Contributing factors that perpetuate ineffective patient-provider communication include the lack of a systematic method for nursing assessment, evaluation, and monitoring of patient-provider communication needs and interventions

and a lack of standardized training of healthcare providers. We propose a call to action for nursing administrators to position patient-provider communication as a patient safety-care quality priority within the healthcare organization and incorporate bedside practices that achieve effective patient communication, especially

with those most vulnerable to impaired communication. Effective patient-provider communication is an essential component of patient care, and for communication to be effective, the information must be complete, accurate, timely, unambiguous, and understood by the patient.¹ By formally implementing the assessment of

Authors' Affiliations: Anesthesiology Resident, University of Michigan, Ann Arbor (Dr Patak); Director, the Joint Commission, Chicago, Illinois (Ms Wilson-Stronks); Speech Language Pathologist, Children's Hospital Boston, Massachusetts (Mr Costello); Director and Professor, Rush University Medical Center and Our Lady of the Resurrection Medical Center, Chicago, Illinois (Dr Kleinpell); Assistant Professor, University of Massachusetts-Amherst School of Nursing (Dr Henneman); Vice President, Creative Health Care Management, Minneapolis, Minnesota (Ms Person); Assistant Professor, University of Pittsburgh School of Nursing, Pennsylvania (Dr Happ).

Disclosure: Dr Patak is the President and CEO for Vidatak, LLC, a for-profit company that focuses on research and development for evidence-based patient communication products and resources.

Mrs Wilson-Stronks is principal investigator for a national study funded by the California Endowment titled, "Hospitals, Language, and Culture: A Snapshot of the Nation," a cross-sectional qualitative study investigating how hospitals are addressing the healthcare needs of diverse patient populations (grant numbers 20032851 and 20062218). She is also coinvestigator for a project funded by the Commonwealth Fund to develop standards for culturally competent patient-centered care in hospitals (grant number 20080055).

Mr Costello, Drs Kleinpell and Henneman, and Mrs Person have no conflicts of interest.

Dr Happ is principal investigator for a clinical trial funded by the National Institute of Child Health and Human Development, the SPEACS study, testing an educational program, SLP consultation and AAC

strategies (R01-HD043988, "Improving Communication with Nonspeaking ICU Patients"). She is coauthor of a copyrighted educational program for ICU nurses used in the SPEACS study. Her research on patient-caregiver communication is currently supported by a grant from NINR K24-NR010244, "Symptom Management, Patient-Caregiver Communication, and Outcomes in ICU."

Corresponding author: Dr Patak, Department of Anesthesiology, University of Michigan, 1H247 UH, SPC 5048, 1500 East Medical Center Dr, Ann Arbor, MI 48109-5048 (lancepat@med.umich.edu).

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.jonajournal.com).



patient communication needs into routine care, nursing administrators will create a sense of accountability among bedside nurses to meet the needs of patients who are communication vulnerable.

A patient's right to effective patient-provider communication is supported by accreditation standards,² regulatory guidelines,^{3,4} and patient rights declarations.^{5,6} Patients have the right to be informed about the care they receive, make educated decisions about their care, and have the right to be listened to by their providers. However, patient communication needs often go unmet or are addressed inappropriately.⁷⁻¹⁰ In the case of non-English-speaking patients, language access services such as the provision of in-person, telephone, or video interpreters and translated documents are either not available or infrequently used.⁸⁻¹¹ Many healthcare institutions rely on ad hoc interpreters such as family, friends, or administrative and custodial staff to communicate and facilitate patient-provider communication, despite the fact that research has shown that the use of ad hoc interpreters can lead to miscommunication and medical errors.¹²

For critically ill or nonspeaking patients, nonverbal behaviors, such as mouthing words, gestures, and head nods, are the principal means of communication; however, these methods have been shown to be ineffective, fatiguing, and inciting frustration.¹³⁻¹⁸ Often, communication is attempted by simply asking yes/no questions, and more appropriate communication interventions are not used. Limiting the patient's communication to yes/no answers restricts the patient's responses to predict-

able messages only or messages that meet the a priori expectation of the patient's need as determined by the clinician.

The absence of effective patient-provider communication has been cited as a significant factor contributing to adverse outcomes.^{19,20} In a 2007 public policy paper focused on health literacy, the Joint Commission recommended that healthcare organizations "make effective communication an organizational priority to protect the safety of patients" and to "incorporate strategies to address patient's communication needs across the continuum of care."²¹ Effective patient-provider communication is a vital component of this transformation and must be prioritized to improve patient safety.

Call to Action

Conduct an Assessment

Patient communication assessment should include a thorough initial assessment of literacy, linguistic, cultural, behavioral, and physical barriers (eg, patient wears glasses or uses hearing aids) at the point of care. It should also include referrals to communication specialists for selection of appropriate interventions when immediate resources at the point of care fail to achieve effective patient communication.

Evaluate the Intervention

An evaluation of the effectiveness and outcomes of communication interventions will determine whether further interventions are necessary.

Monitor and Document Effective Communication

It is imperative that the effectiveness of communication interven-

tions be monitored, as a decline in patient communication may indicate a change in the patient's health status or suggest that an alternative intervention is needed. There are several methods for documenting communication-related information,²²⁻²⁴ and for systematic implementation, Table 1 presents a sample assessment and documentation tool incorporating a methodological sequence of symptom management. This assessment tool was designed to be incorporated into computerized charting menus to assist nurses in selecting drop-down items that corresponded to their patient communication assessment, intervention, and evaluation process. This process can also be incorporated into multidisciplinary rounds by adding patient communication as a topic within patient daily goal sheets or patient centered care assessment forms that are used during multidisciplinary rounds in the ICU or on the wards. (See Form, Supplemental Digital Content 1, to view the daily patient centered rounds goals form, <http://links.lww.com/JONA/A1>.)

Expanding the Multidisciplinary Team and Making Appropriate Referrals

Readily available resources to aid in communication should be present on all patient care units for managing patient communication needs at the point of care. In the event such resources are ineffective, a referral to communication specialists (speech language pathologists, audiologists, and professional healthcare interpreters) may lead to a more comprehensive assessment where the best feature match to a patient's needs can be determined, especially



Table 1. Patient Communication Assessment Tool²⁹

<p>Baseline Communication Method/Special Needs</p> <ul style="list-style-type: none"> (1) Verbal (2) Writing (pen and paper) (3) Communication board (4) Electronic communication device (5) Speaking valve (6) Gesturing (7) Mouthing/lip reading (8) Hearing aids (9) Glasses (10) Language interpreter needed (11) Family facilitated (12) Sign language/interpreter needed (13) Other (document explanation) <p>Assessment</p> <p>Patient's reported level of distress with communication (scale [0-5])^a</p> <ul style="list-style-type: none"> (0) Not at all (1) A little bit (2) Somewhat (3) Quite a bit (4) Very much (5) No response <p>Current Barriers</p> <ul style="list-style-type: none"> (1) Hostility (2) Withdrawn/depressed (3) Delirium (4) Agitation (5) Confusion (6) Impaired level of consciousness (7) Illiterate (8) Orally intubated (9) Tracheotomy (10) Foreign language (11) Sedated (12) Restrained (13) Surgery (14) History of stroke (15) Weakness (16) Vision impairment (17) Hearing impairment (18) Visitation restrictions (19) Other (document explanation) (20) None 	<p>Interventions at Point of Care</p> <ul style="list-style-type: none"> (1) Comfort measures (2) Music (3) Sitter (4) Communication device (document explanation) (5) Phone (6) Speaking valve (7) Calm spoken voice (8) Give patient time to communicate (9) Released restraints (10) Glasses (11) Hearing aid (12) Call light (13) Interpreter (14) Other (document explanation) <p>Reassessment</p> <p>Patient's reported level of distress with communication (scale [0-5])^a</p> <ul style="list-style-type: none"> (0) Not at all (1) A little bit (2) Somewhat (3) Quite a bit (4) Very much (5) No response <p>Evaluation/Effectiveness</p> <ul style="list-style-type: none"> (1) Patient reports being satisfied (2) Family reports being satisfied (3) Patient reports being unsatisfied (4) Family reports being unsatisfied (5) Patient responds appropriately with intervention (6) Necessary information is obtained from and provided to the patient (7) Patient demonstrates understanding (8) Other (document explanation) <p>Referral</p> <ul style="list-style-type: none"> (1) Yes (document explanation) (2) No
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

^aAdapted from Memorial Symptom Assessment Scale–Short Form with permission.³⁰

when the communication process remains dynamic throughout the nonspeaking condition.²⁵ A comprehensive approach to assessment and feature matching and devising a plan of care by speech language pathologists can be quite complex and reinforces the im-

portance of referrals to specialists when point-of-care resources do not achieve desired goals. (See Table, Supplemental Digital Content 2, which gives an overview of an extensive menu of assessment and intervention features, <http://links.lww.com/JONA/A2>.)

Standardize Training for Healthcare Providers

Patient communication strategies, particularly those used to assess and communicate with communication-vulnerable patients, have historically been neglected in medical and nursing education.²⁶ It is important for



healthcare organizations to provide and elevate training on patient-provider communication as an essential component of staff continuing education and development. Recently, commercially available communication boards have been developed and implemented specifically to facilitate commonly used messages with both critically ill and non-English-speaking patients (available in multiple translations at www.vidatak.com).^{27,28} Physicians, nurses, and therapists from various disciplines and other staff who interact directly with nonspeaking and non-English-speaking patients need to be trained on how to work effectively with these communication aids and with interpreters. In addition, nurses should be trained to be sensitive to signs of communication distress and made aware of the process for obtaining appropriate referrals to communication specialists, such as a speech language pathologist. (For samples of communication boards, see Figures, Supplemental Digital Content 3, which is an illustration of the front of a picture board, <http://links.lww.com/JONA/A3>; Supplemental Digital Content 4, which is an illustration of the back of a picture board, <http://links.lww.com/JONA/A4>; Supplemental Digital Content 5, which is an illustration of front of an English-language picture board, <http://links.lww.com/JONA/A5>; Supplemental Digital Content 6, which is an illustration of the back of an English-language picture board, <http://links.lww.com/JONA/A6>; Supplemental Digital Content 7, which is an illustration of the front of an Arabic-language picture board, <http://links.lww.com/JONA/A7>; and Supplemental Digital Content 8, which is an illus-

tration of the back of an Arabic-language picture board, <http://links.lww.com/JONA/A8>.)

Summary

An assessment of communication needs should be done for every patient to determine if patients are able to communicate effectively with healthcare providers or require an intervention (ie, communication boards or other audio or visual aids or interpreters). This should be followed by monitoring for changes in the patient's assessment or changes in the effectiveness of the intervention. The interdisciplinary team should consult with professionals who are trained in specific communication interventions. Healthcare organizations need to have supportive systems in place to help meet patient communication needs, and accreditation and regulatory bodies need to increase attention to this important safety issue as a means to inspire organizations to act. Nursing administrators play a key role in helping to ensure that assessment of communication needs is an integral component of patient care. Improving communication can enhance patient safety, and nurses can serve to champion initiatives to promote patient-provider communication and make a difference in patient outcomes.

Acknowledgments

We thank the people who have made significant contributions, including Christina L. Cordero, PhD, MPH (manuscript preparation and editing), April L. Kopp, MSW, MFA (manuscript preparation and editing), and Bryan Traughber, MD (editing).

Presentations

American Speech and Hearing Association Annual Conference. "AAC and

Communication Vulnerable Patients: A Call to Action;" November 21, 2008; Chicago, Illinois.

The Joint Commission. "Call to Action: Improving Care to Communication Vulnerable Patients." February 17, 2009. Webinar.

REFERENCES

1. Wilson-Stronks A, Lee KK, Cordero CL, et al. *One Size Does Not Fit All: Meeting the Health Care Needs of Diverse Populations*. Oakbrook Terrace, IL: The Joint Commission; 2008.
2. Joint Commission. *The Joint Commission Standards for Hospitals*. Oakbrook Terrace, IL: The Joint Commission; 2008.
3. US Department of Health and Human Services, Office for Civil Rights. Guidance to federal financial assistance recipients regarding title VI prohibition against national origin discrimination affecting limited English proficient persons. 2003. Washington, DC. Available at <http://www.usdoj.gov/crt/cor/lep/hhsrevisedlepguidance.pdf>. Accessed April 29, 2009.
4. US Department of Justice, Civil Rights Division, Disability Rights Section. A guide to disability rights laws. 2005. Available at <http://www.usdoj.gov/crt/ada/cguide.htm>. Accessed April 29, 2009.
5. American Hospital Association. The patient care partnership: understanding expectations, rights, and responsibilities. 2003. Available at http://www.aha.org/aha/content/2003/pdf/pcp_english_030730.pdf. Accessed April 29, 2009.
6. President's Advisory Committee on Consumer Protection and Quality in the Health Care Industry. Final report: quality first: better care for all Americans. 2008. Available at <http://www.hcqualitycommission.gov/final/>. Accessed April 29, 2009.
7. Garrett PW, Dickson HG, Whelan AK. Communication and healthcare complexity in people with little or no English: the Communication Complexity Score. *Ethn Health*. 2008;13(3):203-217.
8. Ramirez D, Engel KG, Tang TS. Language interpreter utilization in the emergency department setting: a clinical review. *J Health Care Poor Underserved*. 2008;19(2):352-362.



9. Regenstein M, Mead H, Muessig KE, et al. Challenges in language services: identifying and responding to patients' needs. *J Immigr Minor Health*. June 7, 2008. Available at http://www.speakingtogether.org/media/file/Challenges_in_Language_Services.pdf. Accessed April 29, 2009.
10. Taveras EM, Flores G. Why culture and language matter: the clinical consequences of providing culturally and linguistically appropriate services to children in the emergency department. *Clin Pediatr Emerg Med*. 2004;5:76-84.
11. Kuo DZ, O'Connor KG, Flores G, et al. Pediatricians' use of language services for families with limited English proficiency. *Pediatrics*. 2007; 119:e920-7.
12. Wilson-Stronks A, Galvez E. *Hospitals, Language, and Culture: A Snapshot of the Nation. Exploring Cultural and Linguistic Services in the Nation's Hospitals: A Report of Findings*. Oakbrook Terrace, IL: The Joint Commission; 2007.
13. Menzel LK. Factors related to the emotional responses of intubated patients to being unable to speak. *Heart Lung*. 1998;27:245-252.
14. Ashworth P. *Care to Communicate. RCN Research Series*. London: Whitefriars Press; 1980.
15. Connolly MA, Shekelton ME. Communicating with ventilator dependent patients. *Dimens Crit Care Nurs*. 1991; 10:115-122.
16. Leathart AJ. Communication and socialisation (1): an exploratory study and explanation for nurse-patient communication in an ITU. *Intensive Crit Care Nurs*. 1994;10:93-104.
17. Happ MB. Communicating with mechanically ventilated patients in ICU: state of the science. *AACN Clin Issues*. 2001;12(2):247-258.
18. Carroll SM. Nonvocal ventilated patients perceptions of being understood. *West J Nurs Res*. 2004;26(1): 85-103; discussion 104-112.
19. Bartlett G, Blais R, Tamblyn R, et al. Impact of patient communication problems on the risk of preventable adverse events in the acute care settings. *CAMJ*. 2008;178:1555-1562.
20. Martinez EL. Patient-centered communication with vulnerable populations. Promising practices for addressing health literacy. Institute of Medicine—Roundtable on Health Literacy; March 29, 2007; Washington, DC.
21. Grubbs V, Chen AH, Bindman AB, et al. Effect of awareness of language law on language access in the health care setting. *J Gen Intern Med*. 2006; 21:683-688.
22. Tatsuya M, Tsunoda J, Inoue S, et al. Communication capacity scale and agitation distress scale to measure the severity of delirium in terminally ill cancer patients: a validation study. *J Palliat Med*. 2001;15:197-206.
23. Williams ML. An algorithm for selecting a communication technique with intubated patients. *Dimens Crit Care Nurs*. 1992;11:222-229.
24. Beukelman DR, Garrett KL, Yorkston KM. *Augmentative Communication Strategies for Adults With Acute Chronic Medical Conditions*. Baltimore, MD: Paul H Brookes Publishing Co; 2007.
25. Costello J. Augmentative communication in the intensive care unit: the Children's Hospital Boston model. *Augment Altern Commun*. 2000;16: 137-153.
26. Shaw J, Hemming MP, Hobson JD, et al. Comprehension of therapy by non-English speaking hospital patients. *Med J Aust*. 1977;2:423-427.
27. Happ MB, Roesch TK, Garrett K. Electronic voice output communication aids for temporarily nonspeaking patients in a medical intensive care unit. *Heart Lung*. 2004;33(2): 92-101.
28. Patak L, Gawlinski A, Fung NI, et al. Communication boards in critical care: patient's views. *Appl Nurs Res*. 2006; 19:182-190.
29. Patak L. *Patient Communication Assessment Tool* [developed at the University of California, Los Angeles Medical Center, Cardiothoracic Intensive Care Unit]. February 2008.
30. Chang VT, Hwang SS, Feuerman M, et al. The Memorial Symptom Assessment Scale Short Form (MSAS-SF) validity and reliability. *Cancer*. 2000; 89(5):1162-1171.